

Tagungsbeitrag zu: Workshop Kommission III „Bodenökologie und Bodenbiologie“ der Deutschen Bodenkundlichen Gesellschaft „Experimenting with Earthworms“  
Veranstalter: Kom. III der DBG  
20.-21. März 2009 in Trier  
Berichte der DBG (nicht begutachtete online Publikation) <http://www.dbges.de>

## Functional importance of earthworm burrowing in field depending on soil tillage

Stefan Schrader<sup>1</sup>

Keywords: Earthworm, burrow, tillage, soil

### Functions of earthworm burrow systems

Primary functions: (1) habitat for earthworms (2) protection from predation (3) protection from UV-radiation (4) space for reproduction and development (5) maintenance of relatively constant conditions.

Secondary functions: (1) gas transport and ventilation (2) water transport and infiltration (3) transport of soluble compounds (4) reduction of erosion (5) improvement of nutrient availability (6) improvement of root growth (7) habitat for other organisms.

### Earthworms controlled by tillage systems in arable soil

One central demand in sustainable agriculture is to bring soil use and soil protection in line (SCHRADER 2001). Reduced tillage systems like conservation tillage and direct drill (no-till) improve biological activity due to less mechanical impact and higher amounts of crop residues on or near the soil surface compared to conventional tillage with ploughing.

Field studies revealed significant higher earthworm cast production and more burrow volume and total length of burrows under conservation compared to conventional tillage which is related to higher biomass

and abundances of earthworms (LANGMAACK et al. 1999; SCHRADER & LARINK 2003). A significant higher abundance of the anecic earthworm species *L. terrestris* caused an improved hydraulic conductivity under direct drill and conservation tillage (BOLL et al. 2006). Detailed species analysis related to population sizes in arable soil revealed a significant higher cast production of the endogeic *A. caliginosa* compared to *L. terrestris* (SCHRADER & LARINK 2003).

### Outcome and conclusions

Reduced tillage systems promote earthworms resulting in

- higher soil turnover
- more functional macropores
- improved hydraulic conductivity

Main drivers of lumbricogenic soil structure development are

- endogeic earthworm species for aggregate formation
- anecic earthworm species for continuous macropore formation

### Literature

- Boll A, Schrader S & Brunotte J 2006. Einfluss von Regenwürmern auf Infiltrationseigenschaften eines Ackerbodens bei unterschiedlicher Bodenbearbeitung. Oral presentation at the conference of Kom. III and VIII of the DBG in Braunschweig 27-29 Sep. 2006
- LANGMAACK M, SCHRADER S, RAPP-BERNHARDT U & KOTZKE K 1999. Quantitative analysis of earthworm burrow systems with respect to biological soil-structure regeneration after soil compaction. *Biology and Fertility of Soils* 28, 219-229.
- SCHRADER S 2001. Ackerboden als Nutz- und Schutzgut aus ökologischer Sicht. *Wasser und Boden* 53, 4-8.
- SCHRADER S & LARINK O 2003. Earthworms as promoters of soil structure rehabilitation. *Tearman: Irish Journal of Agri-Environmental Research* 3, 47-55.

<sup>1</sup> vTI, Institut für Biodiversität, Bundesallee 50, D-38116 Braunschweig  
E-mail: [stefan.schrader@vti.bund.de](mailto:stefan.schrader@vti.bund.de)